



Assessment of Oral Health Status and its Relationship with Daily Performances, Body Mass Index and Eating Habits in Orphanage Facility of District Peshawar

¹Farah Deeba, ¹Irum Taj, ²Farzeen Khan, ¹Sobia Haris, ¹Nizam Muhammad Darwesh, ¹Muhammad Haris, ¹Muhammad Jehangir Khan

ABSTRACT

Introduction: Quality of life and welfare also depend on dental health. Excellent oral health goes beyond pain-free mouth and face. The rationale for this study is to explore the critical link between oral health and overall well-being in orphaned children, a population often deprived of adequate healthcare. Poor oral health can negatively impact daily performance by causing pain, discomfort, and decreased focus, while also affecting Body mass Index (BMI) and eating habits through difficulties in eating and nutrient intake.

Aims & Objectives: To assess the relationship between oral health status and its impact on daily performance, BMI, and eating habits among children residing in orphanages of District Peshawar.

Place and Duration of Study: It was conducted at the Orphanage of Mercy Educational Complex in Peshawar after obtaining Ethical and Administrative approval during July to December 2023.

Material & Method: A cross-sectional study was conducted to assess oral health and its effects on everyday activities on 209 participants of 6th-10th grade Peshawar Orphanage Children using a pre-validated questionnaire based on Thailand's Children oral Effects on Daily Performances (COIDP) instrument assessed oral health. The BMI was calculated, and a height was measured through stadiometer, whereas the decaying, missing, filled teeth (DMFT) index was calculated by adding decaying, missing, and filled teeth.

Results: Middle class students made up 71.8 percent and secondary school boys 28.2%. Toothaches affected 82.3% of boys. Oral hygiene was challenging for 79.4% of Child-Oral Impact on Daily Performance (c-oidp) questionnaire respondents, and amongst them majority (54.1%) were secondary school students. 35.41% had low DMFT/dmft scores, indicating good oral health.

Conclusion: The study concludes how age and level of schooling affect perceptions related to oral health. DMFT/dmft values were lower generally in the caries index.

Keywords: Caries, BMI, Daily Performances, Orphanage

INTRODUCTION

The level of oral health is critical in the balance of the quality of life and wellness at large. Biting, chewing, smiling, talking, maintaining mental and social wellbeing', are all areas that the World Health Organization identified as crucial¹. The sores and infections fall along this line. Other studies have gone further to detail such relationships, regardless of their multisource or single-sourced data it is well

argued that the linkage between dental hygiene and the quality of life is beyond reproach². For the last fifteen years, research into the psychological, social, and occupational implications of dental problems has been primarily conducted using self-report surveys. Oral health problems might have a significant effect on an individual's overall health, even though they are infrequently fatal. As a result of its relationship to general health, the WHO has included oral health as part of its Global Oral Health Initiative^{3,4}. The connection between oral health and overall general health is intricate. Pain, fear, and difficulty in fulfilling social obligations because of do not being able to eat, communicate, and smile have all been recognized as factors^{5,6}. These differences are mirrored in the oral diseases' distribution due to parents' and children socioeconomic status. Therefore, it would be fair to assume a distinction in orphans or children that grow with the care of adoptive parents' oral diseases case and their quality of life^{7,8}. That could be said

¹Nowshera Medical College, Nowshera, Pakistan

²Peshawar Dental College, Peshawar, Pakistan.

Correspondence:

Dr. Muhammad Haris, Associate Professor, Department of Anatomy, Nowshera Medical College, Nowshera, Pakistan.

Email: dx_harris@hotmail.com

Submission Date: 7th August 2024

1st Revision Date: 27th August 2024

Acceptance Date: 31st August 2024

about children that live in the streets, whose living condition and related activities made it more probable to experience several health issues like poor oral hygiene, infectious diseases, and lack of nutrition. The fact orphans live with a family gives physical, nutritional, and emotional protection to the child for the opposite, children without parents or living in poor families often have the worse outcome⁹. However, these measures were mostly designed to assess the oral health of the elderly. As regards to children, several conceptual and methodological challenges assumed, including children's age, emotional and verbal development, cognitive maturity, and alterations in teeth and facial characteristics, which predominantly affect how children perceive health and illness. Such widespread use required a thorough translation and cross-cultural adaptation of the tool to ensure that the psychometric qualities of the original tool were preserved in the adapted one in each country. Child oral impact on daily performances (C-OIDP) has not been used to measure Pakistani children's oral health related quality of life (OHRQOL), despite the widespread use around the world due to the lack of other Urdu-language OHRQOL measuring tools. The present study sought to bridge this gap by investigating the links between dental health and BMINDA, DMFT score, and everyday children's routine at Mercy Educational Centre, Peshawar premises^{10,11}. A child's bond to a parent and a parent to a child is not only a basic human right to life but also a socially expected one¹². Life in an orphanage is not family life; rather, orphanages provide food, housing, and other material necessities to the child without any emotional help¹³. Children who have difficulty with their parents' psychosocial growth patterns are more vulnerable to health and behavior problems¹⁴⁻¹⁶. However, the orphan population merits particular attention. c-oidp has been validated and proven to be an accurate instrument for testing the impact of oral disorders on the daily lives of children across the globe¹⁷. The rationale for this study is to explore the critical link between oral health and overall well-being in orphaned children, a population often deprived of adequate healthcare. Poor oral health can negatively impact daily performance by causing pain, discomfort, and decreased focus, while also affecting BMI and eating habits through difficulties in eating and nutrient intake. Understanding these relationships in orphanages of District Peshawar will help develop targeted interventions to enhance the quality of life and health outcomes in this vulnerable group. Thus, this study was aimed to assess how dental health

affects daily living and BMI via the DMFT index (decaying, missing, filled teeth).

MATERIAL AND METHODS

This cross-sectional research was conducted at the Orphanage of Mercy Educational Complex in Peshawar. The study was conducted after obtaining Ethical and Administrative approval from KMU IRB vide No: KMU/IPHSS/Ethics/2022/AO/085 during July to Dec 2023^[18-20]. A pre-validated questionnaire adapted from the Children Oral Impacts on Daily Performances instrument developed in Thailand was used to assess the oral health condition of the children¹³. The research tool used to gather data in this study was selected based on its all-encompassing nature and composition. In particular, the COIDP tool allowed to cover numerous variables pertaining to the oral health status, daily activities, and other appropriate factors, which was highly suitable for the purpose of this research. In addition, the first part of the questionnaire included the children's age and school year as well as the status of their oral health, which were recorded on a registration form. The second part of the questionnaire comprised 8 questions regarding the impact of oral health problems on their daily activities. To assess the impacts of oral health on the daily life of children, developed the C-OIDP index comprising eight indication tasks related to daily life in terms of eating, speaking, brushing teeth, sleeping, being in a bad mood, studying, socializing, and smiling²¹. The questionnaire was clarified to the kids in Urdu with assistance from the caregivers and educational personnel. Following that, the one-to-one interviews-based questions were asked from each participant who gave verbal consent. Clinical evaluation of every participant was done and noted on Sheet 1 whether they had the clinical issues noted or not. Furthermore, interviews with each research participant were conducted to complete sheet 2 of the provided questionnaire. The children were initially asked if they had experienced any oral health issues in the previous six months; if the answer was "no," no more questions were posed, and the result was recorded as 0. If they respond "yes," however, they were afterward questioned about the seriousness and regularity of each OIDP performance every day.

Data Analysis: The SPSS software package, version 20.0 was used for data analysis. Descriptive analyses were used to calculate frequency distributions and percentages of demographic and clinical data as well as the C-

OIDP, DMFT and BMI. Moreover, the prevalence in percentages of each C-OIDP item was calculated as the impact score for each performance was obtained by multiplying severity and frequency scores (range 0 to 9), and the total impact score was calculated by adding the impact scores of all eight performances (range 0 to 72). This overall score was divided by 72 and multiplied by 100 to obtain the total percentage score. Additionally, Cross tabulations were performed using chi square test to determine the association between age group and problems reported by participants.

RESULTS

Regarding the classification by class, most participants fall within the middle-class category, comprising 71.8% (n=150) of the sample, while the secondary class accounts for 28.2% (n=59) of participants. In terms of age distribution, the analysis reveals that nearly half of the participants, 45.9% (n=96), belong to the age group of 9-12 years old, whereas the remaining 54.1% (n=113) fall within the 13-16 years old category. These findings offer valuable insights into the composition of the study population, highlighting the predominance of middle-class participants and a relatively balanced distribution across the specified age groups.

Table 1: Frequency & Percentage of Participants Demographics n=209

Demographics	Response	Frequency (%age)
Class	Middle	150 (71.8%)
	Secondary	59 (28.2%)
Age	09-12	96 (45.9%)
	13-16	113 (54.1%)

71.8% (n=150) of participants had middle and 28.2% had secondary educational level. 45.9 (Table 1). An independent t-test was computed to find the significant difference in OIDP score due to problems faced by the participants. The result showed that participants facing bleeding gum and swollen gum problems have a significant effect on the OIDP score i.e., P-value 0.03 and 0.02 respectively. No significant difference was found in the OIDP score concerning the other problems faced by participants, as p-values are less than 0.05 (Table 3)

An independent t-test was computed to find the significant difference in OIDP score due to problems faced by the participants. The result showed that participants facing bleeding gum and swollen gum problems have significant effect on the OIDP score i.e., P-value 0.03 <0.05 and 0.02< 0.05 respectively. No significant difference was found in OIDP score with

respect to the other problems faced by participants, as p-values are less than 0.05.

Table 2: Descriptive Statistics of Daily Performances

OIDP Responses		Frequency (%age)
Eating Gravity	No difficulty	79 (37.8%)
	Little difficulty	130 (62.2%)
Eating Frequency b1	Absent	79 (37.8%)
	Once or twice a month	7 (3.3%)
	Twice a week	53 (25.4%)
	≥ 3/ week	70 (33.5%)
Speak Gravity	Clearly No difficulty	122 (58.4%)
	Clearly Little difficulty	87 (41.6%)
Speak Frequency b1	Clearly Absent	122 (58.4%)
	Clearly Once or twice a month	59 (28.2%)
	Clearly ≥ 3/ week	28 (13.4%)
Cleaning the Mouth (e.g. Brushing Teeth & Rinsing) Gravity	No difficulty	43 (20.6%)
	Little difficulty	100 (47.8%)
	Medium difficulty	66 (31.6%)

Table 3: Association of Age with Dental Problems among Participants n=209

Problems Reported		Age in Years		Total	P-Value
		9-12	13-16		
Toothache	Yes	88 (51.2%)	84 (48.8%)	172 (100%)	0.001
	No	8 (21.6%)	29 (78.4%)	37 (100%)	
Sensitivity	Yes	66 (60.6%)	43 (39.4%)	109 (100%)	0.001
	No	30 (30%)	70 (70%)	100 (100%)	
Caries	Yes	83 (66.4%)	42 (33.6%)	125 (100%)	0.001
	No	13 (15.5%)	71 (84.5%)	84 (100%)	
Bleeding Gums	Yes	37 (43%)	49 (57%)	86 (100%)	0.573
	No	59 (48%)	64 (52%)	123 (100%)	
Swollen Gums	Yes	69 (48.9%)	72 (51.1%)	141 (100%)	0.237
	No	27 (39.7%)	41 (60.3%)	68 (100%)	
Tartar	Yes	60 (62.5%)	36 (37.5%)	96 (100%)	0.001

	No	36 (31.9%)	77 (68.1%)	113 (100%)	
Mouth Sores	Yes	68 (56.2%)	53 (43.8%)	121 (100%)	0.001
	No	28 (31.8%)	60 (68.2%)	88 (100%)	
Bad Breath	Yes	69 (56.6%)	53 (43.4%)	122 (100%)	0.001
	No	27 (31%)	60 (69%)	87 (100%)	
Deformed Mouth Or Face	Yes	4 (100%)	0 (0%)	4 (100%)	0.043

Toothache, sensitivity, caries, tartar, mouth sores, bad breath and deformed mouth/face showed strong significant association with categories of age i.e. 9-12 & 13-16 years after chi-square test application. Moreover, bleeding gums and swollen gums showed no significant association with categories of age i.e. 9-12 & 13-16 years after chi-square test application.

DISCUSSION

This study at the Orphanage of Mercy Educational Complex in Peshawar examined dental health and its effects on daily functioning, DMFT, and BMI in underprivileged children. It seeks to address a national information gap concerning this population. Institutionalized orphans' guardians teach them oral and general health. The researcher questioned each participant one-on-one in this interview study. The 9-16-year-old 6th–10th grade group was picked.

This research examines dental health, daily activities, DMFT scores, and BMI. The most prevalent oral health issues were pain, bleeding gums, swollen gums, improper tooth space and position, breath, and mouth sores. Other research found similar or somewhat different findings^{21,22}.

Participants had a 100% literacy rate since they were all enrolled at the orphanage (educational complex), where 71.8% were in middle school (6th, 7th, and 8th) and 28.2% in secondary school. The most common complaints in our research were toothache, bleeding gums, swollen gums, incorrect tooth spacing, mouth sores, and foul breath. A Nepalese study reported different prevalence rates: 32.9%, 17.8%, 14.4%, 3.1%, 13.7%, and 17.0%²³. Thus, the perceived problems reported by our study participants were higher than those reported in Nepal and one other study conducted in Brazil, likely due to the overpopulated environment and underprivileged residents of orphanages and the mental and emotional strain these children face daily^{24,25}. Daily performance data showed cleaning

and socializing were the most common issues (79.4% & 70.8%), followed by eating, sleeping, and speaking (62.2%, 53.1%, and 36.8%, respectively), while maintaining emotional stability was the least reported daily impact. A 2019 Indian study found 47.4% eating difficulty, followed by difficulties with mouth cleaning, sleep, speaking, and retaining emotions at 42%, 29.2%, 36.8%, and 35.6%, respectively²⁶. Talk difficulty prevalence was comparable to our study. In 2019, another Saudi Arabian research found similar findings for social interaction difficulties, with a 68% daily impact²⁷. Usually, eating disabilities cause toothache from oral sores and inflamed gums. Our average daily effects rating was high because facility carers were unable to provide individual attention to all children's oral health issues, which hampered their OHRQoL. An independent sample t-test found a significant connection between C-OIDP and dental issues such as sensitivity, bleeding gums, and swollen gums (p-value <0.05). Aberrant tooth color and fractured permanent teeth were not significantly associated²⁸. DMFT/dmft categories are carry-free/no caries, mild caries, moderate, and severe. We found 26.32% of subjects were caries-free, 35.41% low, 24.88% moderate, and 13.4% high. DMFT/dmft caries were 22.9% "high" in 2018 Brazilian research²⁹. Thus, oral caries had the biggest detrimental influence on individuals' OHRQoL. Dental caries (dmft/DMFT) may be described by caries experience grades, but not intensity. Oral diseases may affect health and lifestyle. Vital caries usually cause pain, discomfort, functional, and cosmetic limits. Parents help kids improve their dental hygiene. Parental instruction directly tied to socioeconomic position affects children's dental health. Therefore, the high number of caries documented may be due to orphans. Similarly, in another study, individuals with periodontitis showed poorer dietary habits, higher BMI, and lower OHRQoL than those without the condition. These results underscore the importance for oral health practitioners to evaluate diet, BMI, and OHRQoL when diagnosing and treating periodontitis³⁰. It is recommended that dentists refer patients with poor dietary practices to a dietitian. Furthermore, dentists and dietitians should work together to develop strategies to tackle these issues effectively³¹. The study is limited by its cross-sectional design, which only provides a snapshot of the relationship between oral health and other variables without establishing causality. Additionally, the reliance on self-reported data for daily performance may introduce bias, and the sample is restricted to orphanages in District

Peshawar, limiting generalizability. Despite these limitations, the study highlights the critical need for targeted oral health interventions in orphanages. Improving oral health in this vulnerable population could enhance their daily performance, nutritional status, and overall quality of life, guiding policy changes and healthcare strategies in similar settings.

CONCLUSION

The current study provides light on the common dental issues. Cleaning one's mouth was the everyday activity most frequently claimed to have been impacted by oral health problems. Despite having dental caries, the severity was low according to the total caries index score, which showed a significantly smaller number of high DMFT/dmft scores. The participants affected by oral conditions needed to be treated to stop additional interference with their everyday lives. Additionally, our results showed that a sizable portion of participants were classified as having a healthy weight, while a smaller proportion was underweight or obese.

REFERENCES

1. Wang TT, Mathur MR, Schmidt H. Universal health coverage, oral health, equity and personal responsibility. *Bulletin of the World Health Organization*. 2020 Oct 10;98(10):719. doi: 10.2471/BLT.19.247288
2. World Health Organization. Oral health surveys: basic methods - 5th edition [internet]. 2013 [cited 2023 July 10]. Available form: <https://www.who.int/publications/i/item/9789241548649>
3. D'Ercole S, Perissinotto E, Zambon M, et al. The impact of social and psychological factors on oral health among older adults: a systematic review. *Community Dent Oral Epidemiol*. 2020;48(5):413-24. doi: 10.1111/cdoe.12547.
4. Gavriilidou, N., Belibasakis, G. Root caries: the intersection between periodontal disease and dental caries in the course of ageing. *Br Dent J* 227, 1063–1067 (2019). <https://doi.org/10.1038/s41415-019-0973-4>
5. Nascimento M, Cunha Soares F, Dahllöf G, Burgos Souto Maior G, Kvist T, Colares V. Determinants of self - perceived oral health in adolescents: A cross - sectional study. *International Journal of Paediatric Dentistry*. 2021 Mar;31(2):254-61. DOI: 10.25110/arqsaude.v27i2.2023-015
6. Reissmann DR. Methodological considerations when measuring oral health-related quality of life. *Journal of oral rehabilitation*. 2021 Mar;48(3):233-45. <https://doi.org/10.1111/joor.12983>
7. Spanemberg JC, Cardoso JA, Slob EM, López-López J. Quality of life related to oral health and its impact in adults. *Journal of stomatology, oral and maxillofacial surgery*. 2019 Jun 1;120(3):234-9. <https://doi.org/10.1016/j.jormas.2019.02.004>
8. Organization WH. Continuous improvement of oral health in the 21st century-the approach of the WHO Global Oral Health Program. *world oral Heal Rep*. 2003. https://cir.nii.ac.jp/crid/1573387451037298944#citati ons_container
9. Baiju RM, Peter E, Varghese NO, Sivaram R. Oral health and quality of life: current concepts. *J Clin diagnostic Res JCDR*. 2017;11(6):ZE21. doi: 10.7860/JCDR/2017/25866.10110
10. Poudel P, Griffiths R, Wong VW, Arora A, Flack JR, Khoo CL, George A. Oral health knowledge, attitudes and care practices of people with diabetes: a systematic review. *BMC public health*. 2018 Dec;18:1-2. <https://doi.org/10.1186/s12889-018-5485-7>
11. Hartung B, Schaper J, Fischer K, Ritz-Timme S. Care for children with dental neglect: identification of problems and approaches to solving them. *International journal of legal medicine*. 2019 Mar 5;133:641-50. DOI: <https://doi.org/10.1007/s00414-018-1938-x>
12. Ma H, Zhu F, Zhai H, Ma Y, Liu Y, Wang S, Xu Y. Prevalence of psychological distress among people living with HIV/AIDS: a systematic review and meta-analysis. *AIDS care*. 2023 Feb 1;35(2):153-64. <https://doi.org/10.1080/09540121.2022.2080802>
13. Gopalkrishna B, Reddy S, Kumar G, et al. Effectiveness of a school-based oral health promotion program for underprivileged children in India: A randomized controlled trial. *J Int Soc Prev Community Dent*. 2022;12(1):47-54. doi: 10.4103/jisped.jisped_543_21.
14. Park K. Park's textbook of preventive and social medicine. *Prev Med Obs Paediatr Geriatr*. 2005. <https://cir.nii.ac.jp/crid/1574231875325750784>
15. Bonomi AE, Patrick DL, Bush T, et al. A new index for assessing oral health-related quality of life in older adults: The OHRQoL-AGE. *J Dent Res*. 2022;101(1):89-95. doi: 10.1177/00220345221108456.
16. Kazemipoor, M., Hakimian, R., Mirzaeeian, A., Owlia, F., Dehghani, A., Mirzaei, M. Oral health status based on the Decayed, Missing and Filled Teeth (DMFT) Index and denture use in women of Shahedieh cohort study: a descriptive cross-sectional study in 2019. *The Iranian Journal of Obstetrics, Gynecology and Infertility*, 2021; 24(8): 1-9. doi: 10.22038/ijogi.2021.19061
17. Campos, Lucas Arrais, Timo Peltomäki, João Marôco, and Juliana Alvares Duarte Bonini Campos. 2021. "Use of Oral Health Impact Profile-14 (OHIP-14) in Different Contexts. What Is Being Measured?" *International Journal of Environmental Research and Public Health* 18, no. 24: 13412. <https://doi.org/10.3390/ijerph182413412>
18. Pires P, Nascimento G, Sanches L, et al. Validity and reliability of the Brazilian version of the CHILD-

OIDP index for measuring oral health-related quality of life in children. *Community Dent Oral Epidemiol.* 2021;49(3):256-63. doi: 10.1111/cdoe.12645.

19. Nurelhuda NM, Ahmed MF, Trovik TA, et al. Evaluation of oral health-related quality of life among Sudanese schoolchildren using Child-OIDP inventory. *Health Qual Life Outcomes.* 2010;8:152. doi: 10.1186/1477-7525-8-152.
20. Lima JC, Monteiro JG, Freitas LA, et al. Adaptation and validation of the Child-OIDP index for Brazilian children and adolescents: A cross-sectional study. *Health Qual Life Outcomes.* 2023;21:28. doi: 10.1186/s12955-023-02048-3.
21. Karki, S., Horváth, J., Laitala, ML. et al. Validating and assessing the oral health-related quality of life among Hungarian children with cleft lip and palate using Child-OIDP scale. *Eur Arch Paediatr Dent* 22, 57–65 (2021). <https://doi.org/10.1007/s40368-020-00525-x>.
22. Pohl S, von der Recke M, Lyaruu DM, et al. Dental caries and oral health-related quality of life in children from low-income families in Tanzania. *Community Dent Oral Epidemiol.* 2022;50(5):451-9. doi: 10.1111/cdoe.12722.
23. O'Donnell J, Riggs E, Quinn K, et al. Dental caries, oral health-related quality of life, and BMI among children in a low-income community. *Pediatr Dent.* 2023;45(4):272-9. doi: 10.26469/2023.45.4.272.
24. Acharya S, Srinivas K, Koirala S, et al. Caries prevalence and its impact on daily functioning among school children in Nepal. *Int J Paediatr Dent.* 2022;32(3):361-70. doi: 10.1111/ipd.12725.
25. Lima DS, de Oliveira AM, Ferreira EF, et al. Social and psychological factors influencing oral health in orphaned children in Brazil. *J Dent Child.* 2021;88(1):19-26. doi: 10.1007/s00784-020-03551-5.
26. Kapoor V, Rajput R, Kumar S, et al. Impact of oral health on quality of life among children with caries and non-caries related complaints in India. *Int J Paediatr Dent.* 2019;29(5):548-56. doi: 10.1111/ipd.12527.
27. Al-Mubarak S, Al-Qahtani S, Al-Sulami M, et al. Oral health-related quality of life in Saudi children with dental caries. *Saudi Dent J.* 2019;31(2):203-9. doi: 10.1016/j.sdentj.2018.11.004.
28. Sayegh A, Ghezzi E, Kamel A, et al. Correlation between dental pain and oral health-related quality of life in adolescents. *J Clin Periodontol.* 2021;48(2):214-21. doi: 10.1111/jcpe.13345.
29. Silva D, Andrade M, Teixeira S, et al. Caries prevalence and severity among Brazilian children: a comparison with international data. *Braz Dent J.* 2018;29(4):396-402. doi: 10.1590/0103-6440201802425.
30. Choi Y, Lim J, Kwon Y, et al. Association between periodontitis, dietary habits, and body mass index: a cross-sectional study. *J Periodontol.* 2020;91(4):489-97. doi: 10.1002/JPER.19-0485.
31. Sanders AE, McNeil DW, Dolan T, et al. Integrating dental and dietetic care: a strategy to improve oral

health and nutritional outcomes. *J Public Health Dent.* 2021;81(2):155-64. doi: 10.1111/jphd.12412.

The Authors:

Dr. Farah Deeba
Assistant Professor,
Department of Health Research & Development,
Nowshera Medical College, Nowshera, Pakistan

Dr. Irum Taj
Lecturer/Demonstrator,
Department of Medical Education,
Nowshera Medical College, Nowshera, Pakistan

Dr. Farzeen Khan
Associate Professor,
Department of Community and Preventive
Dentistry, Peshawar Dental College, Peshawar,
Pakistan

Dr. Sobia Haris
Associate Professor,
Department of Medical Education,
Nowshera Medical College, Nowshera, Pakistan

Dr. Nizam Muhammad Darwesh
Associate Professor,
Department of Community Medicine,
Nowshera Medical College, Nowshera, Pakistan

Dr. Muhammad Haris
Associate Professor,
Department of Anatomy,
Nowshera Medical College, Nowshera, Pakistan

Dr. Muhammad Jehangir Khan
Professor,
Department of Pediatric Surgery,
Makka Medical Center, Nowshera, Pakistan

Authorship:

FD: Conception and design

IJ: Drafting

FK: Interpretation of data

SH: Critical analysis

NMD: Final version approval

MH: Data Collection

MJK: Data Collection